

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use BUPROPION HYDROCHLORIDE EXTENDED-RELEASE TABLETS (XL) safely and effectively. See full prescribing information for BUPROPION HYDROCHLORIDE EXTENDED-RELEASE TABLETS (XL).

BUPROPION HYDROCHLORIDE extended-release tablets USP (XL) for oral use
Initial U.S. Approval: 1985

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

See full prescribing information for complete boxed warning.

- Increased risk of suicidal thinking and behavior in children, adolescents, and young adults taking antidepressants. Monitor for worsening and emergence of suicidal thoughts and behaviors. (5.1)

RECENT MAJOR CHANGES

Boxed Warning 05/2017
Warnings and Precautions, Neuropsychiatric Adverse Events and Suicide Risk in Smoking Cessation (5.2) 05/2017

INDICATIONS AND USAGE

Bupropion hydrochloride extended-release tablets (XL) are amine oxidase antidepressant, indicated for the treatment of major depressive disorder (MDD) and prevention of seasonal affective disorder (SAD). Periodically reevaluate long-term usefulness for the individual patient. (1)

DOSEAGE AND ADMINISTRATION

- Increase dose gradually to reduce seizure risk. (2.1, 5.3)
- Periodically reassess the dose and need for maintenance treatment. (2.2)

Major Depressive Disorder:

- Starting dose: 150 mg once daily. Usual target dose: 300 mg once daily (2,2)
 - After 4 days, may increase the dose to 300 mg once daily. (2,2)
- Seasonal Affective Disorder:**
- Initiate treatment in the autumn prior to onset of seasonal depressive symptoms. (2,3)
 - Starting dose: 150 mg once daily. Usual target dose: 300 mg once daily. (2,3)
 - After one week, may increase the dose to 300 mg once daily. (2,3)
 - Continue treatment through the winter season. (2,3)

Heptic Impairment:

- Moderate to severe hepatic impairment: 150 mg every other day (2,6)
- Mild hepatic impairment: Consider reducing the dose and/or frequency of dosing. (2, 8, 7)

Renal Impairment:

- Consider reducing the dose and/or frequency of dosing. (2, 7, 8, 6)

ADVERSE REACTIONS AND STRENGTHS

Extended-release tablets: 150 mg, 300 mg (3)

CONTRAINDICATIONS

- Seizure disorder. (4, 5, 3)
- Current or prior diagnosis of bulimia or anorexia nervosa. (4, 5, 3)
- Avoid discontinuation of alcohol, benzodiazepines, barbiturates, antiepileptic drugs. (4, 5, 3)
- Monamine oxidase inhibitors (MAOIs): Do not use MAOIs intended to treat psychiatric disorders with bupropion hydrochloride extended-release tablets (XL) within 14 days of stopping treatment with bupropion hydrochloride extended-release tablets (XL). Do not use bupropion hydrochloride extended-release tablets (XL) within 14 days of stopping an MAOI intended to treat psychiatric disorders. In addition, do not start bupropion hydrochloride extended-release tablets (XL) in a patient who is being treated with linezolid or intravenous methylene blue. (4, 7, 6)
- Known hypersensitivity to bupropion or other ingredients of bupropion hydrochloride extended-release tablets (XL). (4, 5, 8)

- Neuropsychiatric Adverse Events: Smoking Cessation: Postmarketing reports of serious or clinically significant neuropsychiatric adverse events have included changes in mood (including depression and mania), psychosis, hallucinations, paranoia, delusions, homicidal ideation, aggression, hostility, aggression, anxiety, and panic, as well as suicidal ideation, suicide attempt, and completed suicide. Observe patients attempting to quit smoking with bupropion hydrochloride extended-release tablets (XL) for the occurrence of such symptoms and instruct them to discontinue bupropion hydrochloride extended-release tablets (XL) and contact a healthcare provider if they experience such adverse events. (5.2)
- Seizure Risk: The risk is dose-related. Can minimize risk by limiting daily dose to 450 mg and gradually increasing the dose. Discontinue if seizure occurs. (4, 5, 3, 7, 3)
- Hypertension: Bupropion hydrochloride extended-release tablets (XL) can increase blood pressure. Monitor blood pressure before initiating treatment and periodically during treatment. (5.4, 6)
- Activation of Mania/Hypomania: Screen patients for bipolar disorder and monitor for these symptoms. (5.5)
- Paranoia and Other Neuropsychiatric Reactions: Instruct patients to contact a healthcare professional if such reactions occur. (5, 6)
- Angle-Closure Glaucoma: Angle-closure glaucoma has occurred in patients with untreated anatomically narrow angles treated with antidepressants. (5, 7)

- ADVERSE REACTIONS: Most common adverse reactions are incidence $\geq 5\%$; $\geq 2\times$ (placebo rate): dry mouth, nausea, insomnia, dizziness, pharyngitis, abdominal pain, agitation, anxiety, tremor, palpitation, sweating, tinnitus, myalgia, anorexia, urinary frequency, constipation, flatulence, and abdominal pain. (2, 6)

To report SUSPECTED ADVERSE REACTIONS, contact TAPI Pharmaceuticals, Inc. at 1-844-518-2989 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- CYP2D6 inducers: Dose increase may be necessary if administered with CYP2D6 inducers (e.g., ritonavir, lopinavir, efavirenz, carbamazepine, phenobarbital, and phenytoin) based on clinical exposure, but should not exceed the maximum recommended dose. (2, 6)
- Drugs metabolized by CYP2D6: Bupropion inhibits CYP2D6 and can increase concentrations of antidepressants (e.g., venlafaxine, nortriptyline, imipramine, desipramine, paroxetine, fluoxetine, sertraline), antipsychotics (e.g., haloperidol, risperidone, thioridazine), beta-blockers (e.g., metoprolol), and Type 1C antiarrhythmics (e.g., propafenone, flecainide). Consider dose reduction when using with bupropion. (7, 2)
- Drugs that lower seizure threshold: Do not use bupropion hydrochloride extended-release tablets (XL) with tramadol. (5, 2, 7, 3)
- Dopaminergic Drugs (levodopa and amantadine): CNS toxicity can occur when used concomitantly with bupropion hydrochloride extended-release tablets (XL). (7, 4)
- MAOIs: Increased risk of hypertensive reactions can occur when used concomitantly with bupropion hydrochloride extended-release tablets (XL). (7, 6)
- Drug laboratory test interactions: Bupropion hydrochloride extended-release tablets (XL) can cause false-positive urine test results for amphetamines. (7, 2)

USE IN SPECIFIC POPULATIONS

- Pregnancy: Use only if benefit outweighs potential risk to the fetus. (8, 1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 08/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

1 INDICATIONS AND USAGE

1.1 Major Depressive Disorder

1.2 Seasonal Affective Disorder

2 DOSEAGE AND ADMINISTRATION

2.1 General Instructions for Use

2.2 Dosage for Major Depressive Disorder (MDD)

2.3 Dosage for Seasonal Affective Disorder (SAD)

2.4 Switching Patients from WELLBUTRIN Tablets (bupropion hydrochloride tablets) or from WELLBUTRIN SR (bupropion hydrochloride extended-release tablets) (SR) to bupropion hydrochloride extended-release tablets (XL)

2.5 To Discontinue Bupropion Hydrochloride Extended-Release Tablets (XL), Taper the Dose

2.6 Dosage Adjustment in Patients with Hepatic Impairment

2.7 Dose Adjustment in Patients with Renal Impairment

2.8 Switching a Patient to or from a Monamine Oxidase Inhibitor (MAOI) Antidepressant

2.9 Use of Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3 DOSEAGE FORMS AND STRENGTHS

3.1 Bupropion Hydrochloride Extended-Release Tablets (XL)

3.2 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

4 CONTRAINDICATIONS

4.1 Contraindications

4.2 Contraindications

4.3 Contraindications

4.4 Contraindications

4.5 Contraindications

4.6 Contraindications

4.7 Contraindications

4.8 Contraindications

4.9 Contraindications

4.10 Contraindications

4.11 Contraindications

4.12 Contraindications

4.13 Contraindications

4.14 Contraindications

4.15 Contraindications

4.16 Contraindications

4.17 Contraindications

4.18 Contraindications

4.19 Contraindications

4.20 Contraindications

4.21 Contraindications

4.22 Contraindications

4.23 Contraindications

4.24 Contraindications

4.25 Contraindications

4.26 Contraindications

4.27 Contraindications

4.28 Contraindications

4.29 Contraindications

4.30 Contraindications

4.31 Contraindications

4.32 Contraindications

4.33 Contraindications

4.34 Contraindications

4.35 Contraindications

4.36 Contraindications

4.37 Contraindications

4.38 Contraindications

4.39 Contraindications

4.40 Contraindications

4.41 Contraindications

4.42 Contraindications

4.43 Contraindications

4.44 Contraindications

4.45 Contraindications

4.46 Contraindications

4.47 Contraindications

4.48 Contraindications

4.49 Contraindications

4.50 Contraindications

4.51 Contraindications

4.52 Contraindications

4.53 Contraindications

4.54 Contraindications

4.55 Contraindications

FULL PRESCRIBING INFORMATION

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

Antidepressants increased the risk of suicidal thoughts and behavior in children, adolescents, and young adults in short-term trials. These trials did not show an increase in the risk of suicidal thoughts and behavior with antidepressant use in subjects aged 65 and older. *See Warnings and Precautions (5.1).*

In patients of all ages who are started on antidepressant therapy, monitor closely for worsening, and for emergence of suicidal thoughts and behaviors. Advise families and caregivers of the need for close observation and communication with the prescriber. *See Warnings and Precautions (5.1).*

1 INDICATIONS AND USAGE

1.1 Major Depressive Disorder

Bupropion hydrochloride extended-release tablets (XL) are indicated for the treatment of major depressive disorder (MDD), as defined by the Diagnostic and Statistical Manual (DSM).

The efficacy of the immediate-release formulation of bupropion was established in two 4-week controlled inpatient trials and one 6-week controlled outpatient trial of adult patients with MDD. The efficacy of the sustained-release formulation of bupropion hydrochloride extended-release tablets (XL) was established in a long-term (up to 44 weeks), placebo-controlled trial in patients who had responded to bupropion in an 8-week study of acute treatment. *[See Clinical Studies (14.1)].*

1.2 Seasonal Affective Disorder

Bupropion hydrochloride extended-release tablets (XL) are indicated for the prevention of seasonal major depressive episodes in patients with a diagnosis of seasonal affective disorder (SAD). The efficacy of bupropion hydrochloride extended-release tablets (XL) in the prevention of seasonal major depressive episodes was established in 3 placebo-controlled trials in adult outpatients with a history of MDD with an autumn-winter seasonal pattern as defined in the DSM. *[See Clinical Studies (14.2)].*

2 DOSEAGE AND ADMINISTRATION

2.1 General Instructions for Use

To minimize the risk of seizure, increase the dose gradually. *See Warnings and Precautions (5.3).* Bupropion hydrochloride extended-release tablets (XL) should be swallowed whole and not crushed, divided, or chewed.

Bupropion hydrochloride extended-release tablets (XL) should be administered in the morning and may be taken with or without food.

2.2 Dosage for Major Depressive Disorder (MDD)

The recommended starting dose for MDD is 150 mg once daily in the morning. After 4 days of dosing, the dose may be increased to the target dose of 300 mg once daily in the morning.

It is generally agreed that acute episodes of depression require several months or longer of antidepressant treatment beyond the response in the acute phase. It is not known if a longer duration of treatment with bupropion hydrochloride extended-release tablets (XL) does need for maintenance treatment is identical to that provided an initial response. Periodically reassess the need for maintenance treatment and the appropriate dose for such patients.

2.3 Dosage for Seasonal Affective Disorder (SAD)

The recommended starting dose for SAD is 150 mg once daily. After 7 days of dosing, the dose may be increased to the target dose of 300 mg once daily in the morning. Doses above 300 mg of bupropion HCl extended-release tablets are not assessed in the SAD trials.

For the prevention of seasonal MDD episodes associated with SAD, initiate bupropion hydrochloride extended-release tablets (XL) in the autumn, prior to the onset of depressive symptoms. Continue treatment through the winter season. Taper and discontinue bupropion hydrochloride extended-release tablets (XL) in early spring. For patients treated with 300 mg per day, decrease the dose to 150 mg once daily before discontinuing bupropion hydrochloride extended-release tablets (XL). Individualize the timing of initiation, and duration of treatment should be individualized, based on the patient's historical pattern of seasonal MDD episodes.

2.4 Switching Patients from WELLBUTRIN Tablets (bupropion hydrochloride tablets) or from WELLBUTRIN SR (bupropion hydrochloride extended-release tablets) (SR) to bupropion hydrochloride extended-release tablets (XL)

When switching patients from WELLBUTRIN Tablets (bupropion hydrochloride tablets) to bupropion hydrochloride extended-release tablets (XL) or from WELLBUTRIN SR (bupropion hydrochloride extended-release tablets) (SR) to bupropion hydrochloride extended-release tablets (XL), give the same total daily dose.

To discontinue bupropion hydrochloride extended-release tablets (XL), taper the dose when discontinuing treatment in patients treated with bupropion hydrochloride extended-release tablets (XL) 300 mg once daily, decrease the dose to 150 mg once daily prior to discontinuation.

2.5 Dosage Adjustment in Patients with Hepatic Impairment

In patients with moderate to severe hepatic impairment (Child-Pugh score: 7 to 15), the maximum dose is 150 mg every other day. In patients with mild hepatic impairment (Child-Pugh score: 5 to 6), consider reducing the dose and/or frequency of dosing. *See Use in Specific Populations (8.7) and Clinical Pharmacology (12.3).*

2.6 Dose Adjustment in Patients with Renal Impairment

In patients with moderate to severe renal impairment (creatinine clearance ≤ 30 mL/min), consider reducing the dose and/or frequency of dosing. *See Use in Specific Populations (8.7) and Clinical Pharmacology (12.3).*

2.7 Switching a Patient to or from a Monamine Oxidase Inhibitor (MAOI) Antidepressant

Discontinue bupropion hydrochloride extended-release tablets (XL) before initiating treatment and initiation of therapy with bupropion hydrochloride extended-release tablets (XL). Conversely, at least 14 days should be allowed after stopping bupropion hydrochloride extended-release tablets (XL) before starting an MAOI antidepressant (5, 6).

2.8 Use of Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

Do not start bupropion hydrochloride extended-release tablets (XL) in a patient who is being treated with a reversible MAOI such as linezolid or intravenous methylene blue. Drug interactions can increase risk of hypertensive reactions. In a patient who requires more urgent treatment of a psychiatric condition, non-pharmacological interventions, including hospitalization, should be considered. *See Contraindications (4.1).*

In some cases, a patient already receiving therapy with bupropion hydrochloride extended-release tablets (XL) may require urgent treatment with linezolid or intravenous methylene blue. Bupropion hydrochloride extended-release tablets (XL) and intravenous methylene blue treatment are not available and the potential benefits of linezolid or intravenous methylene blue treatment are judged to outweigh the risks of hypertensive reactions in a particular patient. In such cases, the combination of bupropion hydrochloride extended-release tablets (XL) and intravenous methylene blue can be administered. The patient should be monitored for 7 weeks or until 24 hours after the last dose of linezolid or intravenous methylene blue, whichever comes first. Therapy with bupropion hydrochloride extended-release tablets (XL) may be resumed 24 hours after the last dose of linezolid or intravenous methylene blue. *See Contraindications (4.1) and Drug Interactions (7.6).*

2.9 Use of Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

Do not start bupropion hydrochloride extended-release tablets (XL) in a patient who is being treated with a reversible MAOI such as linezolid or intravenous methylene blue. Drug interactions can increase risk of hypertensive reactions. In a patient who requires more urgent treatment of a psychiatric condition, non-pharmacological interventions, including hospitalization, should be considered. *See Contraindications (4.1).*

In some cases, a patient already receiving therapy with bupropion hydrochloride extended-release tablets (XL) may require urgent treatment with linezolid or intravenous methylene blue. Bupropion hydrochloride extended-release tablets (XL) and intravenous methylene blue treatment are not available and the potential benefits of linezolid or intravenous methylene blue treatment are judged to outweigh the risks of hypertensive reactions in a particular patient. In such cases, the combination of bupropion hydrochloride extended-release tablets (XL) and intravenous methylene blue can be administered. The patient should be monitored for 7 weeks or until 24 hours after the last dose of linezolid or intravenous methylene blue, whichever comes first. Therapy with bupropion hydrochloride extended-release tablets (XL) may be resumed 24 hours after the last dose of linezolid or intravenous methylene blue. *See Contraindications (4.1) and Drug Interactions (7.6).*

3 DOSEAGE FORMS AND STRENGTHS

3.1 Bupropion Hydrochloride Extended-Release Tablets (XL)

3.2 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.3 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.4 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.5 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.6 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.7 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.8 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.9 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.10 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.11 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.12 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.13 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.14 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.15 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.16 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.17 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.18 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.19 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.20 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.21 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.22 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.23 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.24 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.25 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.26 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.27 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.28 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.29 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.30 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.31 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.32 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.33 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.34 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.35 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.36 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.37 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.38 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.39 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

3.40 Bupropion Hydrochloride Extended-Release Tablets (XL) with Reversible MAOIs such as Linezolid or Methylene Blue

- Bupropion hydrochloride extended-release tablets (XL) may have an odor. This is normal.

- Take your doses of bupropion hydrochloride extended-release tablets (XL) at least 8 hours apart.

- You may take bupropion hydrochloride extended-release tablets (XL) with or without food.

- It is not dangerous to smoke and take bupropion hydrochloride extended-release tablets (XL) at the same time. But, you will lower your chance of breaking your smoking habit if you smoke after the date you set to stop smoking.

- You may use bupropion hydrochloride extended-release tablets (XL) and nicotine patches (a type of nicotine replacement therapy) at the same time, following the precautions below.

- You should only use bupropion hydrochloride extended-release tablets (XL) and nicotine patches together under the care of your healthcare provider. Using bupropion hydrochloride extended-release tablets (XL) and nicotine patches together may raise your blood pressure, and sometimes this can be severe.
 - You should only use bupropion hydrochloride extended-release tablets (XL) and nicotine patches together under the care of your healthcare provider. Using bupropion hydrochloride extended-release tablets (XL) and nicotine patches together may raise your blood pressure, and sometimes this can be severe.

- Tell your healthcare provider if you plan to use nicotine patches. Your healthcare provider should check your blood pressure regularly if you use nicotine patches with bupropion hydrochloride extended-release tablets (XL) to help you quit smoking.

- If you miss a dose, do not take an extra dose to make up for the dose you missed. Wait and take your next dose at the regular time. **This is very important.** Too much bupropion hydrochloride extended-release tablets (XL) can increase your chance of having a seizure.

- If you take too much bupropion hydrochloride extended-release tablets (XL), or overdose, call your local emergency room or poison control center right away.

Do not take any other medicines while taking bupropion hydrochloride extended-release tablets (XL) unless your healthcare provider has told you it is okay.

What should I avoid while taking bupropion hydrochloride extended-release tablets (XL)?

- Limit or avoid using alcohol during treatment with bupropion hydrochloride extended-release tablets (XL). If you usually drink a lot of alcohol, talk with your healthcare provider before suddenly stopping. If you suddenly stop drinking alcohol, you may increase your chance of having seizures.

- Do not drive a car or use heavy machinery until you know how bupropion hydrochloride extended-release tablets (XL) affect you. Bupropion hydrochloride extended-release tablets (XL) can affect your ability to do these things safely.

What are possible side effects of bupropion hydrochloride extended-release tablets (XL)?

Bupropion hydrochloride extended-release tablets (XL) can cause serious side effects. See the sections at the beginning of this Medication Guide for information about serious side effects of bupropion hydrochloride extended-release tablets (XL).

The most common side effects of bupropion hydrochloride extended-release tablets (XL) include:

- trouble sleeping
- feeling anxious
- stuffy nose
- nausea
- dry mouth
- constipation
- dizziness
- joint aches

If you have trouble sleeping, do not take bupropion hydrochloride extended-release tablets (XL) too close to bedtime.

Tell your healthcare provider right away about any side effects that bother you.

These are not all the possible side effects of bupropion hydrochloride extended-release tablets (XL). For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

You may also report side effects to Twi Pharmaceuticals, Inc. at 1-844-518-2989.

How should I store bupropion hydrochloride extended-release tablets (XL)?

- Store bupropion hydrochloride extended-release tablets (XL) at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

- Protect from light.

Keep bupropion hydrochloride extended-release tablets (XL) and all medicines out of the reach of children.

General information about the safe and effective use of bupropion hydrochloride extended-release tablets (XL).

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use bupropion hydrochloride extended-release tablets (XL) for a condition for which it was not prescribed. Do not give bupropion hydrochloride extended-release tablets (XL) to other people, even if they have the same symptoms you have. It may harm them.

If you take a urine drug screening test, bupropion hydrochloride extended-release tablets (XL) may make the test result positive for amphetamines. If you tell the person giving you the drug screening test that you are taking bupropion hydrochloride extended-release tablets (XL), they can do a more specific drug screening test that should not have this problem.

This Medication Guide summarizes important information about bupropion hydrochloride extended-release tablets (XL). If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about bupropion hydrochloride extended-release tablets (XL) that is written for health professionals.

For more information about bupropion hydrochloride extended-release tablets (XL), go to www.twipharma.com or call 1-844-518-2989.

What are the ingredients in bupropion hydrochloride extended-release tablets (XL)?

Active ingredient: bupropion hydrochloride.

Inactive ingredients: hypromellose, glyceryl behenate, hydroxypropyl cellulose, colloidal silicon dioxide, ethylcellulose, povidone, methacrylic acid copolymer type C, triethyl citrate, iron oxide black and propylene glycol. The tablets are printed with edible black ink.

Manufactured for:

Twi Pharmaceuticals USA, Inc.
Paramus, NJ 07652

Manufactured by:

Twi

Twi Pharmaceuticals, Inc.
Taoyuan City, 32063, Taiwan

WELLBUTRIN XL® is a registered trademark of GlaxoSmithKline LLC used under license.

All other product/brand names are the trademarks of their respective owners.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

20780

Revised: 08/2017

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Bupropion hydrochloride extended-release tablets (XL)

Buprop